Richmond

7110 Forest Ave, Suite 203 Richmond, VA 23226



Prince George

2025 Waterside Rd, Suite 100B Prince George, VA 23875

Physician Order - Remicade (Infliximab)

PATIENT INFORMATION		
Name: DOB:		
Allergies: Phone Number:		
REFERRAL STATUS		
☐ New Referral ☐ Dose/Frequency Change ☐ Order Renewal		
Location Preference (optional)		
☐ Richmond ☐ Prince George		
If patient has a central line, then the placement report, diagnostic imaging to confirm tip placement and date of last access are required		
DIAGNOSIS AND ICD-10 CODE		
☐ Moderate to Severe Ulcerative Colitis	K51.90	
☐ Moderate to Severe Crohn's Disease	K50.90	
Rheumatoid Arthritis	M06.9	
☐ Ankylosing Spondylitis	M45.9	
Psoriatic Arthritis	L40.50	
Plaque Psoriasis	L40.0	
Other:	ICD10	
PEOLINED DOCUMENTATION (must include)		
REQUIRED DOCUMENTATION (must include) ☐ This signed order form by the provider ☐ Clinical/Progress Notes		
_		
☐ Patient Demographics AND Insurance Inform ☐ Negative QuantiFERON Gold TB Test or Skin F		Tests Supporting Primary Dx
☐ Patient currently receiving same therapy at Last dose:		
List Tried & Failed Therapies, including duration of treatment		
1) 3)		
2) 4)		
-7		
MEDICATION ORDERS**		
Initial Dosing Remicade 3mg/kg infuse IV over 2 hours at week 0, 2, 6, then every 8 weeks		
☐ Remicade 5mg/kg infuse IV over 2 hours at week 0, 2, 6, then every 8 weeks		
☐ Remicademg/kg infuse IV over 2 hours at week 0, 2, 6, then every 8 weeks		
Maintenance Dosing Remicade 5mg/kg infuse IV over 2 hours every 8 weeks		
☐ Remicade 10mg/kg infuse IV over 2 hours every 8 weeks		
Alternative Dosing Remicade mg/kg infuse IV over 2 hours every weeks		
Patient Weight: kg or lb Patient Height: in		
Duration: x 6 months x 1 year doses		
**For all weight-based therapies, Infusion Solutions will obtain/verify a patient's current weight within one week of dosing.		
OPTIONAL PREMEDICATIONS and LAB ORDERS		
☐ Acetaminophen 650mg PO prior to infusion		
☐ Diphenhydramine 25mg PO or IV (per patient preference) prior to infusion		
☐ Methylprednisolone 40mg slow IVP prior to infusion		
☐ CMP drawn yearly		
☐ Other PreMed or Lab Order with frequency:		
In the event of an infusion reaction or adverse event, our covering physician will be notified and appropriate medical care will be administered.		
PRESCRIBER INFORMATION		
Prescriber Name:	NPI:	Contact:
Phone: Fax:		Email:
Prescriber Signature:		Date: