Richmond

7110 Forest Ave, Suite 203 Richmond, VA 23226



Prince George

2025 Waterside Rd, Suite 100B Prince George, VA 23875

Physician Order - Renflexis (Infliximab-abda)

PATIENT INFORMATION			
Name: DOB:			
Allergies:		Phone Numbe	er:
·			
REFERRAL STATUS			
☐ New Referral ☐ Dose/Frequency Change ☐ Order Renewal			
Location Preference (optional)			
☐ Richmond ☐ Prince George			
If patient has a central line, then the placement report, diagnostic imaging to confirm tip placement and date of last access are required			
DIAGNOSIS AND ICD-10 CODE			
☐ Moderate to Severe Ulcerative Colitis	K51.90		
☐ Moderate to Severe Crohn's Disease	K50.90		
☐ Rheumatoid Arthritis	M06.9		
Ankylosing Spondylitis	M45.9		
☐ Psoriatic Arthritis	L40.50		
☐ Plaque Psoriasis	L40.0		
☐ Other:	ICD10		
REQUIRED DOCUMENTATION (must include)			
This signed order form by the provide			rogress Notes
			Tests Supporting Primary Dx
Negative QuantiFERON Gold TB Test or Skin PPD HBV Panel results			
Patient currently receiving same therapy at Last dose:			
List Tried & Failed Therapies, including duration of treatment			
1)	3)		
2)	4)		
MEDICATION ODDEDS**			
MEDICATION ORDERS**			
Initial Dosing Renflexis 3mg/kg infuse IV over 2 hours at week 0, 2, 6, then every 8 weeks			
☐ Renflexis 5mg/kg infuse IV over 2 hours at week 0, 2, 6, then every 8 weeks ☐ Renflexis mg/kg infuse IV over 2 hours at week 0, 2, 6, then every 8 weeks			
Maintenance Dosing Renflexis 5mg/kg infuse IV over 2 hours every 8 weeks			
Renflexis 10mg/kg infuse IV over 2 hours every 8 weeks			
Alternative Dosing Renflexis mg/kg infuse IV over 2 hours every weeks			
The make bosing Weeks			
Patient Weight: kg or lb Patient Height: in			
Duration: x 6 months x 1 year doses			
**For all weight-based therapies, Infusion Solutions will obtain/verify a patient's current weight within one week of dosing.			
OPTIONAL PREMEDICATIONS and LAB ORDERS			
Acetaminophen 650mg PO prior to infusion			
☐ Diphenhydramine 25mg PO or IV (per patient preference) prior to infusion			
☐ Methylprednisolone 40mg slow IVP prior to infusion			
☐ CMP drawn yearly			
Other PreMed or Lab Order with frequency:			
In the event of an infusion reaction or adverse event, our covering physician will be notified and appropriate medical care will be administered.			
PRESCRIBER INFORMATION			
Prescriber Name: NPI: Contact:			
Phone:	Fax:		Email:
Prescriber Signature:	1. 4		Date:
O			