IV Iron Therapies
Referring Physician Orders Rev. 3/2023
Contact us with questions at: 804-442-3558 or email: referrals@theinfusionsolution.com
Please fax completed referral form & all required documents to 804-554-5848



|   |   | PATIENT DEMOG  | RAPH   | ICS               |                       |               |                |
|---|---|--|--|-------------------|-----------------------|---------------|----------------|
| Patient Name:   |   |  | DOB: Phone:  |                   |                       |               |                |
| Address:  |   |  | City/ST/Zip:   |                   |                       |               |                |
|   |   |  | IKDA   |                   | □ lbs □ kg            |               |                |
|   |   | INFORMATION: Please attac  |  |                   | _                     |               |                |
| DIAGNOSIS*  |   |  |  |                   |                       |               |                |
| *ICD 10 Code  | ☐ Iron Deficiency Anemia (D5                            |  |  | Other:            |                       | ICI           | 240            |
| Required  | E non Bendency Anemia (Bo                               | ,,   |  |                   |                       | , ici         | 510            |
|   |   | INFUSION C   |  |                   |                       |               |                |
| MEDICATION DOSE  Feraheme® (ferumoxytol) 510 mg   |   |  | DIRECTIONS/DURATION  |                   |                       |               |                |
| Feraheme® (ferumoxyto   |   | ☐ Two-dose regimen: Infuse IV over 30 minutes once, followed by a second IV dose over 30 minutes 3-8 days later. |  |                   |                       |               |                |
|   |   |  | *Administer while patient is in reclined or semi-reclined position.*   |                   |                       |               |                |
|   |   |  |  |                   | utes after completion |               |                |
| INFed® (iron dextran)   | ☐ 100mg<br>☐mg  |  | <ul> <li>□ TEST Dose: 25mg IV over 30 seconds x 1 dose. Observe for 1 hour before administering remainder of therapeutic dose.</li> <li>□ Infuse 100mg IV over 60 minutes every day for a total of doses.</li> <li>□ Infuse IV over hours (range 4-6 hours) x 1 dose.</li> </ul> |                   |                       |               |                |
|   | *Maximum dos  | □ lof  |  |                   |                       |               |                |
|   |   | □ Infu   |  |                   |                       |               |                |
| Injectafer® (ferric carbox  | ymaltose) TWO-dose regime<br>□ ≥50kg: 750m              | _ LIW  | ☐ TWO-dose regimen: Infuse IV over 15-30 minutes once, followed by a second IV dose over 15-30 minutes 7 days later.   |                   |                       |               |                |
|   | •   | <sup>9</sup>   |  |                   |                       |               |                |
|   | ONE-dose regimen  | <u>1</u>   | ☐ ONE-dose regimen: Infuse IV over 15-30 minutes x 1 dose.   |                   |                       |               |                |
|   | □ ≥50kg:<br>*Maximum dos                                | mg (15 mg/kg)  |  |                   |                       |               |                |
| Monoferric® (ferric derise  |   | -  | ☐ Infuse IV over 15-30 minutes x 1 dose.   |                   |                       |               |                |
| Wichordino (Terrio denoc  |   | mg (20 mg/kg)  | LI IIIIUSE IV UVEI 13-30 IIIIIIIUIES X I UUSE.   |                   |                       |               |                |
| Venofer® (iron sucrose)   | □ 100mg   | □ Info   | use 100m   | g IV over 30 minu | utes 3 times weekly x | doses.        |                |
| □ 200mg   |   |  |  |                   | utes 3 times weekly x | doses.        |                |
|   | receiving therapy above fr                              | -  |  |                   | <b>5</b>              |               |                |
| If yes, Facility Nam  | e:  |  |  |                   | Date of n             | ext treatment |                |
| OTHER ORDERS  |   |  |  |                   |                       |               |                |
| Labs to be drawn by: ☐ Infusion Center ☐ Referring Physician  |   |  |  |                   |                       |               |                |
| ☐ No labs ordered   | at this time<br>□ CMP q                                 | □ CRP a □ F  | SR a   | П                 | l FTs a               | □ Other:      |                |
| PRE-MEDICATION O  | ·   |  |  |                   |                       |               |                |
| ☐ No premeds ordered at this time   |   |  | □ Diphe  | nhydramine 2      | 5mg PO                |               |                |
| ☐ Acetaminophen 650mg PO  |   |  | ☐ Methy  | Iprednisolone     | 40mg IVP -OR-         | ☐ Hydrocortis | sone 100mg IV  |
| ☐ Other:  |   |  |  |                   |                       |               |                |
|   |   | REFERRING PHYSICIA   | AN INF   | ORMATION          |                       |               |                |
| Physician Signature:  |   |  |  |                   | Date:                 |               |                |
| Physician Name:   |   | Provider NPI:  |  |                   | Specialty:            |               |                |
| Address:  |   |  | City/ST/Zip:   |                   |                       |               |                |
| Contact Person: Phone #:  |   | Phone #:   |  |                   | Fax #:                |               |                |
| Email Where Follow Up   | Documentation Should Be Se                              | ent:   |  |                   |                       |               |                |
|   | F   | REQUIRED CLINICAL [  | OOCUN  | MENTATION         |                       |               |                |
| Please attach i   | medical records: Initial H&I                            |  |  |                   |                       | ults to suppo | ort diagnosis  |
|   | iron deficiency anemia confi                            |  | 7.00, mo   | aloution hot, t   | and labortest res     | uito to ouppt | ort diagnosis. |
| ☐ Yes ☐ No Does the patient require intravenous iron therapy for iron supplementation? If yes, please select all that app |   |  |  |                   |                       |               |                |
|   | ☐ Inadequate response to p☐ Contraindication to oral th |  | <ul> <li>☐ Intolerance to prior oral therapy</li> <li>☐ Decreased absorption of oral iron</li> </ul>   |                   |                       |               |                |
|   | (e.g., inflammatory bowel                               |  |  |                   | ing gastric bypass    |               |                |
|   | ☐ Iron (blood) loss at rate to                          | ate too rapid for oral intake to   Other:  |  |                   |                       |               |                |
| LAD AND TEST SEC  | compensate for the loss                                 |  |  |                   |                       |               |                |
| LAB AND TEST RES  |   | elitin   |  |                   |                       |               |                |
|   | erum Iron   |  |  |                   |                       |               |                |
| PRIOR FAILED THERAPIES (oral/IV iron supplementation)  Medication Failed: Dates of Treatment: Reason for D/C:             |   |  |  |                   |                       |               |                |
|   |   |  |  |                   |                       |               |                |
|   |   |  |  |                   |                       |               |                |
|   |   | = 2.00 0. 1100.110111.   |  |                   |                       |               |                |